

***BVRI* Observations of the Optical Afterglow of GRB 990510¹**

K. Z. Stanek² and P. M. Garnavich

Harvard-Smithsonian Center for Astrophysics, 60 Garden St., MS20, Cambridge, MA 02138

e-mail: kstanek@cfa.harvard.edu, peterg@cfa.harvard.edu

J. Kaluzny

Warsaw University Observatory, Al. Ujazdowskie 4, PL-00-478 Warszawa, Poland

and

N. Copernicus Astronomical Center, Bartycka 18, PL-00-716 Warszawa, Poland

e-mail: jka@sirius.astrouw.edu.pl

W. Pych

Warsaw University Observatory, Al. Ujazdowskie 4, PL-00-478 Warszawa, Poland

e-mail: pych@sirius.astrouw.edu.pl

I. Thompson

Carnegie Observatories, 813 Santa Barbara St., Pasadena, CA 91101-1292

e-mail: ian@ociw.edu

ABSTRACT

We present *BVRI* observations of the optical counterpart to the Gamma-Ray Burst (GRB) 990510 obtained with the Las Campanas 1.0-m telescope between 15 and 48 hours after the burst. The temporal analysis of the data indicates steepening decay, independent of wavelength, approaching asymptotically $t^{-0.76 \pm 0.01}$ at early times ($t \ll 1$ day) and $t^{-2.40 \pm 0.02}$ at late times, with the break time at $t_0 = 1.57 \pm 0.03$ days. GRB 990510 is the most rapidly fading of the well-documented GRB afterglows. It is also the first observed example of broad-band break for a GRB optical counterpart. The optical spectral energy distribution, corrected for significant Galactic reddening, is well fitted by a single power-law with $\nu^{-0.61 \pm 0.12}$. However, when the *B*-band point is dropped from the fit, the power-law becomes $\nu^{-0.46 \pm 0.08}$, indicating a possible deviation from the power-law in the spectrum, either intrinsic or due to additional extinction near the source or from an intervening galaxy at $z = 1.62$. Broad-band break behavior broadly similar to that observed in GRB 990510 has been predicted in some jet models of GRB afterglows, thus supporting the idea that the GRB energy is beamed, at least in some cases.

Subject headings: gamma-rays: bursts

¹Based on the observations collected at the Las Campanas Observatory 1 m Swope telescope

²On leave from N. Copernicus Astronomical Center, Bartycka 18, Warszawa PL-00-716, Poland

1. INTRODUCTION

The BeppoSAX satellite (Boella et al. 1997) has brought a new dimension to gamma-ray burst (GRB) research, by rapidly providing good localization of several bursts per year. This has allowed many GRBs to be followed up at other wavelengths, including X-ray (Costa et al. 1997), optical (van Paradijs et al. 1997) and radio (Frail et al. 1997). Good positions have also allowed redshifts to be measured for a number of GRBs (e.g. GRB 970508: Metzger et al. 1997), providing a definite proof for their cosmological origin.

GRB 990510 was detected by BeppoSAX (Piro et al. 1999) on May 10.36743 UT, and the fluence was among the highest of the BeppoSAX localized events, after GRB 990123, GRB 980329, and GRB 970111 (Amati et al. 1999). It was also detected by BATSE (Kippen et al. 1999) and its peak flux and fluence rank it in the top 4% (9%) of BATSE burst flux (fluence) distributions. The BeppoSAX NFI follow-up of GRB 990510 started about 8 hours after the burst (Kuulkers et al. 1999) and detected a strong X-ray afterglow.

Axelrod, Mould & Schmidt (1999) were the first to optically monitor the field of GRB 990510, starting on 10.514 UT, i.e. only 3.5 *hours* after the burst, using the Mount Stromlo 50-inch telescope. The optical counterpart to GRB 990510 was first identified by Vreeswijk et al. (1999a) with data taken about 8.5 *hours* (May 10.73 UT) after the burst, using the Sutherland 1.0-m telescope. It was easily recognized as a bright ($R \approx 19.2$; but see below), new object not present in the DSS image at the position of $\alpha = 13^h 38^m 07^s.64$, $\delta = -80^\circ 29' 48''.8$ (J2000.0) (Kaluzny et al. 1999b). Galama et al. (1999) confirmed the optical transient (OT) with the ESO 2.2-m telescope on May 10.99 UT and found it to be declining with a temporal decay index of -0.85 . Kaluzny et al. (1999a,b) observed the GRB 990510 with the Las Campanas 1.0-m telescope, beginning on May 10.995, and found that the object declined by 0.34 mag in the *R*-band over 4.6 *hours*. They also noticed that the *R*-band reference star used by Vreeswijk et al. (1999) and Galama et al. (1999) is in fact brighter than they assumed by 0.7 mag (USNO-A2.0 catalog: Monet et al. 1996), making the afterglow as bright as $R \approx 17.75$ when first observed by Axelrod et al. (1999). This change in calibration was confirmed by independent photometric calibration of the GRB field by Pietrzyński & Udalski (1999a) and Bloom et al. (1999). Absorption lines at $z = 1.619$ seen in the optical spectrum of GRB 990510 taken with the VLT-UT1 8-m telescope by Vreeswijk et al. (1999b) provide a lower limit to the redshift of the GRB source.

By adding data taken during their second night (starting May 12.0 UT), Stanek et al. (1999) noticed the steepening of the OT decline, and they derived a power-law decay index of -1.36 by a fit to their data from both nights, but they also showed that a single power-law does not provide a good fit to all the data. Analysis of more data has confirmed this behavior (Hjorth et al. 1999; Bloom et al. 1999; Marconi et al. 1999a). We will discuss in detail the temporal behavior of the GRB 990510 OT further in this paper.

We describe the data and the reduction procedure in Section 2. In Section 3 we discuss the multiband temporal behavior of the GRB OT. In Section 4 we describe the broad-band spectral




Fig. 1.— Finding chart for the field of GRB 990510. The optical transient is indicated with an arrow. Also shown are the three comparison stars calibrated by Pietrzyński & Udalski (1999a,b) and Bloom et al. (1999).

properties of the afterglow deduced from our optical data.

2. THE DATA AND THE REDUCTION PROCEDURE

The data were obtained with the Las Campanas Observatory 1.0-meter Swope telescope on two nights, May 10/11 and May 11/12 UT, 1999. We used the backside-illuminated SITE3 2048×4096 CCD, with the pixel scale of $0.44 \text{ arcsec pixel}^{-1}$. To speed up the readout time we used only a 2048×1200 section of the CCD. We have obtained 35 *R*-band, 20 *V*-band, 16 *I*-band and 4 *B*-band images, with exposure times ranging from 240 *sec* during the first night to 900 *sec* during the second night³ The image quality was good, with median seeing for the *R*-band of $1.4''$.

³*BVRI* data are available through anonymous ftp on cfa-ftp.harvard.edu, in `pub/kstanek/GRB990510` directory, and through the WWW at <http://cfa-www.harvard.edu/cfa/oir/Research/GRB/>.

The finding chart for the GRB 990510 is shown in Figure 1.

The data were reduced using two different methods. First, we have used the photometric data pipeline of the the DIRECT project (Kaluzny et al. 1998; Stanek et al. 1998), which is based on the Daophot PSF-fitting image reduction package (Stetson 1987; 1992). To check the consistency of the differential photometry we have also employed the image subtraction method of Alard & Lupton (1999) and Alard (1999a), as implemented in the ISIS image subtraction package (Alard 1999b). We have found a very good agreement between these two data reduction methods (~ 0.01 mag).

The calibration for the field has been obtained by Pietrzyński & Udalski (1999a,b) in the *BVI* bands and by Bloom et al. (1999) in the *VR* bands. In both cases the quoted uncertainty of the calibration zero point is ± 0.03 mag for the *VRI* bands and ± 0.05 mag for the *B*-band. Comparison of the *V*-band measurements for three secondary standards given by Pietrzyński & Udalski (1999a,b) and Bloom et al. (1999) in the GRB field yields a systematic difference of 0.025 mag, i.e. within the quoted uncertainty. We have adopted the calibration of Pietrzyński & Udalski (1999a,b) for the *BVI* bands and the calibration of Bloom et al. (1999) for the *R*-band.

3. THE TEMPORAL BEHAVIOR

We plot the GRB 990510 *BVRI* lightcurves in Figure 2. Most of the data comes from our monitoring (Kaluzny et al. 1999a,b; Stanek et al. 1999) and from the OGLE project (Pietrzyński & Udalski 1999a,b,c). We use those OGLE points which extend the time coverage beyond that of our data. In addition, we use the early *R*-band data of Axelrod et al. (1999) and Galama et al. (1999) and later *R*-band data points of Marconi et al. (1999a,b), and one late *V*-band point obtained by Beuermann, Reinsch & Hessman (1999). All these data extend beyond our time coverage and are therefore useful for constraining the time evolution of the afterglow.

As noticed by Stanek et al. (1999) using data from two days only, the optical *R*-band data already showed clear departure from the initial shallow power-law of about -0.9 determined by Galama et al. (1999) and Kaluzny et al. (1999a,b) from the first day data. This trend of steepening decay was confirmed later by Hjorth et al. (1999) and others.

To describe the temporal evolution of the GRB 990510 optical counterpart, we use the combined data described earlier and fit to each *VRI* band separately the following four parameters formula (similar to that of Bloom et al. 1999 and Marconi et al. 1999a):

$$F_{\nu}(t) = \frac{k_{\nu} \left(\frac{t}{t_0} \right)^{-a_1}}{1 + \left(\frac{t}{t_0} \right)^{(a_2 - a_1)}}, \quad (1)$$

where k_{ν} is a normalization constant specific for each band such that the magnitude is given




Fig. 2.— *BVRI* light curves of GRB 990510. Our data is shown with circles and OGLE data with squares. Other data used to constrain the fits is shown with triangles (for references see text). Also shown is the simple analytical fit discussed in the text.

by $m(t) = -2.5 \log(F_\nu)$. This formula describes power-law t^{-a_2} decline at early times ($t \ll t_0$) and another power-law t^{-a_1} decline at late times ($t \gg t_0$). The total number of points used for the fits was 44 for the *R*-band, 32 for the *V*-band, 31 for the *I*-band and five for the *B*-band, total of 112 points. The *R*-band data has the most extensive temporal coverage, especially early after the burst. The other bands are not as well constrained, at least with data at hand, which demonstrates the importance of getting multi-band follow-up images as early as possible. The parameters are correlated with each other, so we first fit the *R*-band and then use the derived t_0 value as fixed for the *VI* bands. We also run a combined fit to all *BVRI* data. The results of this combined fit are shown as the continuous lines in the Figure 2 and the best-fit parameters for all the fits are presented in Table 1. The errors shown are based on conditional probability distributions, fixing two of the parameters at their most probable values.

The combined fit does a good job of representing the overall temporal behavior of the *BVRI* data set, but it is clearly dominated by the *R*-band data. Considering that these data represent an inhomogeneous data set, the χ^2/DOF values for the *VI* bands are rather good. The χ^2/DOF is somewhat higher for the *R*-band. As our *R*-band data has somewhat better photometric accuracy than the *VI* data, this might indicate that we see some small, short-timescales departures from the analytical fit. Our frequent observations of the GRB optical counterpart during the first night were obtained to test for such rapid variations. After we subtract the long-term trend derived above, we get a root-mean-square (*rms*) scatter of 0.02 mag, with the largest deviation of 0.08 mag. Comparing to constant field stars with similar magnitude, the observed scatter is consistent with photometric noise. Clearly, brightness variations on time scales from 0.1 to 2 *hours* are small. We note that Hjorth et al. (1999) obtained similar *rms* scatter in their 31 300 *sec* *R*-band images, obtained with the 1.54-m telescope on La Silla at times overlapping with our observations. By combining these two data sets one might be able to search with more sensitivity for correlated variations in the brightness of the afterglow.

4. REDDENING AND BROAD-BAND SPECTRAL ENERGY DISTRIBUTION

The GRB 990510 is located at Galactic coordinates of $l = 304^\circ 94' 26''$, $b = -17^\circ 80' 79''$. To remove the effects of the Galactic interstellar extinction we used the reddening map of Schlegel, Finkbeiner & Davis (1998, hereafter: SFD). As noticed by Stanek (1999), the expected Galactic reddening towards the burst is significant, $E(B - V) = 0.203$, which corresponds to expected values of Galactic extinction $A_B = 0.88$, $A_V = 0.66$, $A_R = 0.53$ and $A_I = 0.40$, for the Landolt (1992) CTIO filters and standard reddening curve of Cardelli, Clayton & Mathis (1989).

We synthesize the *BVRI* spectrum from our data by interpolating the magnitudes to a common time. As discussed in the previous section, the optical colors of the GRB 990510 counterpart do not show significant variation. We therefore select an epoch of May 11.26 UT

Table 1. GRB 990510 BEST-FIT PARAMETERS

Band	a_1	a_2	t_0 (days)	χ^2/DOF
<i>R</i>	2.41 ± 0.02	0.76 ± 0.01	1.55 ± 0.03	1.51
<i>V</i>	2.46 ± 0.05	0.73 ± 0.02	...	1.22
<i>I</i>	2.17 ± 0.07	0.82 ± 0.04	...	1.03
<i>BVRI</i>	2.40 ± 0.02	0.76 ± 0.01	1.57 ± 0.03	1.48




Fig. 3.— Synthetic spectrum of GRB 990510 21.5 *hours* after the burst, constructed using analytical fit shown in Figure 2.

(21.5 *hours* after the burst) for the color analysis, which coincides with our first *B*-band images. Using the all-band fit from the previous section we get for this epoch $B_{21.5} = 20.39 \pm 0.05$, $V_{21.5} = 19.82 \pm 0.03$, $R_{21.4} = 19.40 \pm 0.03$ and $I_{21.5} = 18.93 \pm 0.03$, where the 0.03 mag error bar represents the combined uncertainty in the zero points calibration and the small measurement errors. Using the SFD extinction values, this translates to unreddened values of $B_{0,21.5} = 19.51 \pm 0.10$, $V_{0,21.5} = 19.16 \pm 0.07$, $R_{0,21.5} = 18.87 \pm 0.06$ and $I_{0,21.5} = 18.53 \pm 0.05$ (following SFD we took the error in $E(B - V)$ to be 0.02 mag).

We convert the *BVRI* magnitudes to fluxes using the effective wavelengths and normalizations of Fukugita, Shimasaku & Ichikawa (1995). These conversions are accurate to about 5%, which increases the error-bars correspondingly. Note that while the error in the $E(B - V)$ reddening value has not been applied to the error-bars of individual points, we include it in the error budget of the fitted slope. The results are plotted in Figure 3 for both the observed and the dereddened

magnitudes. The spectrum is well fitted by a single power-law with $\nu^{-0.61 \pm 0.12}$, although when the B -band point is dropped from the fit, the power-law becomes $\nu^{-0.46 \pm 0.08}$, indicating a possible deviation from the power-law in the blue part of the spectrum, either intrinsic or due to additional extinction outside our Galaxy.

5. CONCLUSIONS

We presented *BVRI* observations of the GRB 990510. Our analysis of the data indicates steepening decay, independent of optical wavelength, with power-law behavior $t^{-0.76 \pm 0.01}$ at early times ($t \ll 1$ day) and second power-law $t^{-2.40 \pm 0.02}$ at late times, with the break time at $t_0 = 1.57 \pm 0.03$ days. This is the first observed example of such broad-band break for a GRB OT and it is very well documented thanks to a concerted effort of the astronomical community. We would like to stress the importance of multiband optical observations for the GRB studies, especially early after the burst.

The optical spectral energy distribution, corrected for significant Galactic reddening, is well fitted by a single power-law with $\nu^{-0.61 \pm 0.12}$. However, when the B -band point is dropped from the fit, the power-law becomes $\nu^{-0.46 \pm 0.08}$, indicating a possible deviation from the power-law in the spectrum, either intrinsic or due to additional extinction near the host or at the intervening galaxy at $z = 1.62$.

The temporal behavior of GRB 990510 is broadly similar to predictions of GRB jet models (Rhoads 1999). A break in the light curve is expected when the jet makes the transition to sideways expansion after the relativistic Lorentz factor drops below the inverse of the opening angle of the initial beam. After the break, the temporal power-law index is expected to approach the electron energy distribution index with values between 2.0 and 2.5, which is consistent with the late-time decline observed here for GRB 990510 (Halpern et al. 1999; Sari, Piran & Halpern 1999). However, the early-time decay index and the spectral slope of GRB 990510 are not as well explained by the models. GRB 990510 provides the first direct evidence that, in some cases, GRB energy is not ejected isotropically.

S. Barthelmy, the organizer of the GRB Coordinates Network (GCN), is recognized for his extremely useful effort. The OGLE collaboration is thanked for making their data promptly available to the astronomical community. We thank C. Alard for his support with the ISIS image subtraction package. We thank B. Paczyński and L. A. Phillips for useful discussions and comments on the paper. KZS was supported by the Harvard-Smithsonian Center for Astrophysics Fellowship. JK was supported by NSF grant AST-9528096 to Bohdan Paczyński and by the Polish KBN grant 2P03D011.12. WP was supported by the Polish KBN grant 2P03D010.15

REFERENCES

- Alard, C., & Lupton, R. H. 1998, *ApJ*, 503, 325
- Alard, C. 1999a, *A&A*, in press (astro-ph/9903111)
- Alard, C. 1999b, private communication (<http://www.iap.fr/users/alard/package.html>)
- Amati, L., et al. 1999, *GCN Circ.* 317
- Axelrod, T., Mould, J., & Schmidt, B. 1999, *GCN Circ.* 315
- Beuermann, K., Reinsch, K., & Hessman, F. V. 1999, *GCN Circ.* 331
- Bloom, J. S., et al. 1999, *GCN Circ.*
- Boella, G., et al. 1997, *A&AS*, 122, 299
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- Frail, D. A., Kulkarni, S. R., Nicastro, L., Feroci, M., & Taylor, G. B. 1997, *Nature*, 389, 261
- Galama, T. J., et al. 1999, *GCN Circ.* 313
- Halpern, J. P., Kemp, J., Piran, T., & Bershadsky, M. A. 1999, *ApJ*, 517, L105
- Hjorth, J., et al. 1999, *GCN Circ.* 320
- Kaluzny, J., Stanek, K. Z., Krockenberger, M., Sasselov, D. D., Tonry, J. L., & Mateo, M. 1998, *AJ*, 115, 1016
- Kaluzny, J., Garnavich, P. M., Stanek, K. Z., Pych, W., & Thompson, I. 1999a, *GCN Circ.* 314
- Kaluzny, J., Garnavich, P. M., Stanek, K. Z., Pych, W., & Thompson, I. 1999b, *IAU Circ.* 7164
- Kippen, R. M., et al. 1999, *GCN Circ.* 322
- Landolt, A. 1992, *AJ*, 104, 340
- Marconi, G., et al. 1999a, *GCN Circ.* 329
- Marconi, G., et al. 1999b, *GCN Circ.* 332
- Metzger, M. R., et al. 1997, *Nature*, 387, 879
- Monet, D., et al. 1996, *USNO-SA2.0*, (U.S. Naval Observatory, Washington DC)
- Pietrzyński, G., & Udalski, A. 1999a, *GCN Circ.* 316
- Pietrzyński, G., & Udalski, A. 1999b, *GCN Circ.* 319
- Pietrzyński, G., & Udalski, A. 1999c, *GCN Circ.* 328
- Piro, L., et al. 1999, *GCN Circ.* 304
- Rhoads, J. E. 1999, *ApJ*, submitted (astro-ph/9903399)
- Sari, R., Piran, R., & Halpern, J. P. 1999, *ApJ*, submitted (astro-ph/9903339)
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Stanek, K. Z. 1999, *GCN Circ.* 312

- Stanek, K. Z., Kaluzny, J., Krockenberger, M., Sasselov, D. D., Tonry, J. L., & Mateo, M. 1998, *AJ*, 115, 1894
- Stanek, K. Z., Garnavich, P. M., Kaluzny, J., Pych, W., & Thompson, I. 1999, *GCN Circ.* 318
- Stetson, P. B. 1987, *PASP*, 99 191
- Stetson, P. B. 1992, in *ASP Conf. Ser. 25, Astrophysical Data Analysis Software and Systems I*, ed. D. M. Worrall, C. Bimesderfer, & J. Barnes (San Francisco: ASP), 297
- van Paradijs, J., et al. 1997, *Nature*, 386, 686
- Vreeswijk, P. M., et al. 1999a, *GCN Circ.* 310
- Vreeswijk, P. M., et al. 1999b, *GCN Circ.* 324

Table 2. GRB 990510 *B*-BAND LIGHTCURVE

HJD –2450000	<i>B</i>	σ_B	Hours after the burst
1309.7494	20.345	0.018	21.0636
1309.7914	20.422	0.021	22.0716
1310.5116	21.325	0.062	39.3564
1310.5230	21.446	0.040	39.6300

Table 3. GRB 990510 *V*-BAND LIGHTCURVE

HJD –2450000	<i>V</i>	σ_V	Hours after the burst
1309.5163	19.414	0.020	15.4692
1309.5216	19.404	0.018	15.5964
1309.5525	19.486	0.022	16.3380
1309.5857	19.517	0.020	17.1348
1309.6152	19.559	0.022	17.8428
1309.6387	19.648	0.019	18.4068
1309.6621	19.637	0.021	18.9684
1309.6852	19.703	0.020	19.5228
1309.7086	19.707	0.025	20.0844
1309.7551	19.856	0.026	21.2004
1309.7822	19.844	0.023	21.8508
1309.8096	19.901	0.034	22.5084
1309.8303	19.866	0.029	23.0052
1309.8511	19.929	0.032	23.5044
1309.8716	20.050	0.039	23.9964
1310.4680	20.762	0.059	38.3100
1310.4763	20.769	0.045	38.5092
1310.6795	20.962	0.051	43.3860
1310.6885	20.999	0.052	43.6020
1310.8748	21.194	0.099	48.0732

Table 4. GRB 990510 *R*-BAND LIGHTCURVE

HJD –2450000	<i>R</i>	σ_R	Hours after the burst
1309.4975	18.949	0.022	15.0180
1309.5030	18.962	0.015	15.1500
1309.5255	19.018	0.018	15.6900
1309.5309	19.031	0.017	15.8196
1309.5582	19.038	0.015	16.4748
1309.5637	19.064	0.017	16.6068
1309.5756	19.096	0.015	16.8924
1309.5806	19.115	0.015	17.0124
1309.5909	19.133	0.014	17.2596
1309.6042	19.125	0.017	17.5788
1309.6094	19.169	0.015	17.7036
1309.6204	19.168	0.015	17.9676
1309.6336	19.170	0.013	18.2844
1309.6439	19.200	0.016	18.5316
1309.6570	19.226	0.021	18.8460
1309.6671	19.259	0.018	19.0884
1309.6801	19.250	0.019	19.4004
1309.6905	19.256	0.019	19.6500
1309.7035	19.287	0.021	19.9620
1309.7203	19.326	0.019	20.3652
1309.7362	19.357	0.015	20.7468
1309.7603	19.417	0.021	21.3252
1309.7770	19.389	0.022	21.7260
1309.8044	19.423	0.020	22.3836
1309.8246	19.447	0.020	22.8684
1309.8453	19.535	0.019	23.3652
1309.8659	19.554	0.020	23.8596
1309.8774	19.501	0.020	24.1356
1309.8844	19.633	0.026	24.3036
1310.4841	20.393	0.032	38.6964
1310.4923	20.356	0.032	38.8932
1310.6613	20.516	0.029	42.9492
1310.6695	20.505	0.025	43.1460
1310.8502	20.645	0.040	47.4828
1310.8618	20.681	0.052	47.7612

Table 5. GRB 990510 *I*-BAND LIGHTCURVE

HJD –2450000	<i>I</i>	σ_I	Hours after the burst
1309.5383	18.592	0.026	15.9972
1309.5472	18.570	0.019	16.2108
1309.5705	18.621	0.021	16.7700
1309.5989	18.640	0.018	17.4516
1309.6285	18.757	0.024	18.1620
1309.6519	18.753	0.017	18.7236
1309.6750	18.804	0.024	19.2780
1309.6984	18.834	0.021	19.8396
1309.7307	18.862	0.021	20.6148
1309.7689	18.935	0.020	21.5316
1309.7994	19.021	0.025	22.2636
1309.8188	19.012	0.025	22.7292
1309.8395	19.051	0.026	23.2260
1309.8602	19.060	0.035	23.7228
1310.5375	19.931	0.064	39.9780
1310.5490	19.878	0.080	40.2540